

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1639MLS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS	1		Web Page URLs for STN Seminar Schedule - N. America
NEWS	2		"Ask CAS" for self-help around the clock
NEWS	3	JAN 27	Source of Registration (SR) information in REGISTRY updated and searchable
NEWS	4	JAN 27	A new search aid, the Company Name Thesaurus, available in CA/CAPLUS
NEWS	5	FEB 05	German (DE) application and patent publication number format changes
NEWS	6	MAR 03	MEDLINE and LMEADLINE reloaded
NEWS	7	MAR 03	MEDLINE file segment of TOXCENTER reloaded
NEWS	8	MAR 03	FRANCEPAT now available on STN
NEWS	9	MAR 29	Pharmaceutical Substances (PS) now available on STN
NEWS	10	MAR 29	WPIFV now available on STN
NEWS	11	MAR 29	New monthly current-awareness alert (SDI) frequency in RAPRA
NEWS	12	APR 26	PROMT: New display field available
NEWS	13	APR 26	IFIPAT/IFIUDB/IFICDB: New super search and display field available
NEWS	14	APR 26	LITALERT now available on STN
NEWS	15	APR 27	NLDB: New search and display fields available
NEWS	16	May 10	PROUSDDR now available on STN
NEWS	17	May 19	PROUSDDR: One FREE connect hour, per account, in both May and June 2004
NEWS	18	May 12	EXTEND option available in structure searching
NEWS	19	May 12	Polymer links for the POLYLINK command completed in REGISTRY
NEWS	20	May 17	FRFULL now available on STN
NEWS	21	May 27	STN User Update to be held June 7 and June 8 at the SLA 2004 Conference
NEWS	22	May 27	New UPM (Update Code Maximum) field for more efficient patent SDIs in CAPLUS
NEWS	23	May 27	CAPLUS super roles and document types searchable in REGISTRY
NEWS	24	May 27	Explore APOLLIT with free connect time in June 2004
NEWS EXPRESS			MARCH 31 CURRENT WINDOWS VERSION IS V7.00A, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 26 APRIL 2004
NEWS HOURS			STN Operating Hours Plus Help Desk Availability
NEWS INTER			General Internet Information
NEWS LOGIN			Welcome Banner and News Items
NEWS PHONE			Direct Dial and Telecommunication Network Access to STN
NEWS WWW			CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may

result in loss of user privileges and other penalties.

* * * * * STN Columbus * * * * *

FILE 'HOME' ENTERED AT 14:14:05 ON 04 JUN 2004

=> d his

(FILE 'HOME' ENTERED AT 14:14:05 ON 04 JUN 2004)

=> file medline biosis embase
COST IN U.S. DOLLARS

SINCE FILE ENTRY	TOTAL SESSION
0.21	0.21

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 14:14:32 ON 04 JUN 2004

FILE 'BIOSIS' ENTERED AT 14:14:32 ON 04 JUN 2004
COPYRIGHT (C) 2004 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'EMBASE' ENTERED AT 14:14:32 ON 04 JUN 2004
COPYRIGHT (C) 2004 Elsevier Inc. All rights reserved.

=> s hyperkines? and hypokines?
L1 278 HYPERKINES? AND HYPOKINES?

=> s l1 and pd<1998
'1998' NOT A VALID FIELD CODE
2 FILES SEARCHED...
L2 133 L1 AND PD<1998

=> s l1 and py<1998
2 FILES SEARCHED...
L3 183 L1 AND PY<1998

=> dup rem
ENTER L# LIST OR (END):L3
PROCESSING COMPLETED FOR L3
L4 113 DUP REM L3 (70 DUPLICATES REMOVED)

=> t ti l4 1-20

L4 ANSWER 1 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.
on STN

TI A case of chorea-acanthocytosis successfully treated with posteroventral
pallidotomy.

L4 ANSWER 2 OF 113 MEDLINE on STN
TI Maternally reported fetal activity levels and developmental diagnoses.

L4 ANSWER 3 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.
on STN
TI [A classification of dysarthria and its treatment].
DE LA CLASSIFICATION DES DYSARTHRIES AUX CONCEPTIONS THERAPEUTIQUES.

L4 ANSWER 4 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI Clinical research and analysis of 56 patients with oesophageal motility
disorders.

L4 ANSWER 5 OF 113 MEDLINE on STN
TI The effect of normalization in reducing variability in regional wall
thickening.

L4 ANSWER 6 OF 113 MEDLINE on STN DUPLICATE 1

TI Catatonia and the neuroleptics: psychobiologic significance of remote and recent findings.

L4 ANSWER 7 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.
on STN

TI Pathophysiology of involuntary movements in adults.

L4 ANSWER 8 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

TI Sympathetic activation significance for biliary duct affection development and possible ways of its correction.

L4 ANSWER 9 OF 113 MEDLINE on STN DUPLICATE 2

TI A study of cognitive development and behavior problems in mentally retarded children.

L4 ANSWER 10 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN

TI Pathophysiology and therapy of movement disorders.

L4 ANSWER 11 OF 113 MEDLINE on STN DUPLICATE 3

TI **Hypokinesia** in Huntington's disease.

L4 ANSWER 12 OF 113 MEDLINE on STN DUPLICATE 4

TI Subjective experience of treatment, side-effects, mental state and quality of life in chronic schizophrenic out-patients treated with depot neuroleptics.

L4 ANSWER 13 OF 113 MEDLINE on STN DUPLICATE 5

TI Heterogeneity of left ventricular regional wall thickening following dobutamine infusion in normal human subjects.

L4 ANSWER 14 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN

TI The role of the basal ganglia in movement control.

L4 ANSWER 15 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN

TI Effects of morphine on EEG in rats and their possible relations to hypo- and **hyperkinesia**.

L4 ANSWER 16 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

TI Usefulness of ATP-atropine echocardiography for diagnosis of myocardial ischemia.

L4 ANSWER 17 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
DUPLICATE 6

TI Effects of electrically-stimulated exercise and passive motion on echocardiographically-derived wall motion and cardiodynamic function in tetraplegic persons.

L4 ANSWER 18 OF 113 MEDLINE on STN DUPLICATE 7

TI The diagnostic value of exercise echocardiography in ischemic heart disease in relation to quantitative coronary arteriography.

L4 ANSWER 19 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

TI Locomotor effects of (D-Trp-11)neurotensin and dopamine transmission in rats.

L4 ANSWER 20 OF 113 MEDLINE on STN DUPLICATE 8

TI [Satellite myocytes in postnatal ontogeny and during skeletal muscle adaptation to hyper- and **hypokinesia**].
Miosatelitotsyty v postnatal'nomu ontogenezi i pry adaptatsii skeletnykh m'iaziv do hiper- i hipokinezii.

=> t ti l4 21-40

- L4 ANSWER 21 OF 113 MEDLINE on STN
TI Facial hyper- and **hypokinesia** following face lift surgery.
- L4 ANSWER 22 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI Overview of basal ganglia function.
- L4 ANSWER 23 OF 113 MEDLINE on STN DUPLICATE 9
TI Time course of left ventricular function and coronary patency after saruplase vs streptokinase in acute myocardial infarction. The PRIMI Trial Study Group.
- L4 ANSWER 24 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN
TI Management of Bell's palsy.
- L4 ANSWER 25 OF 113 MEDLINE on STN DUPLICATE 10
TI [The effect of peridural analgesia on uterine contractions].
Effet de l'analgesie peridurale sur la contraction uterine.
- L4 ANSWER 26 OF 113 MEDLINE on STN DUPLICATE 11
TI Changes in myocardial contraction patterns in response to regional ischemia and sympathetic nerve stimulation.
- L4 ANSWER 27 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI Phenomenology and psychopathology of the catatonic syndrome.
- L4 ANSWER 28 OF 113 MEDLINE on STN DUPLICATE 12
TI Clinical applications of coronary sinus retroperfusion during high risk percutaneous transluminal coronary angioplasty.
- L4 ANSWER 29 OF 113 MEDLINE on STN
TI [The role of biliary dyskinesia in the mechanism of the damage to the protective properties of the mucosal barrier in peptic ulcer].
Rol' diskinezii zhelchevyvodiashchikh putei v mekhanizme povrezhdeniia zashchitnykh svoistv slizistogo bar'era pri iazvennoi bolezni.
- L4 ANSWER 30 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN
TI Free radical therapy for a patient with motor dysfunction (I).
- L4 ANSWER 31 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN
TI Zuclopenthixol and thioridazine in the treatment of aggressive, elderly patients: A double-blind, controlled, multicentre study.
- L4 ANSWER 32 OF 113 MEDLINE on STN DUPLICATE 13
TI Biotelemetric investigation of morphine's thermic and kinetic effects in rats.
- L4 ANSWER 33 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN
TI Decreased metabolic rate as an acrolein resistance mechanism in Drosophila melanogaster.
- L4 ANSWER 34 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN
TI Biliary tract motor dysfunction.
- L4 ANSWER 35 OF 113 MEDLINE on STN DUPLICATE 14
TI Intrapataminal infusion of nerve growth factor to support adrenal medullary autografts in Parkinson's disease. One-year follow-up of first

clinical trial.

L4 ANSWER 36 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
DUPLICATE 15

TI SIGNIFICANCE OF RIGHTWARD AXIS SHIFT IN ANTERIOR ACUTE MYOCARDIAL
INFARCTION.

L4 ANSWER 37 OF 113 MEDLINE on STN

TI [Several indicators of protein and nucleic acid metabolism in lymphoid
organs of rats exposed to **hypokinesia** and vitamin B1
deficiency].

Nekotorye pokazateli metabolizma belkov i nukleinovyx kislot v
limfoidnykh organakh kryt v usloviyakh gipokinezii i defitsita vitamina
B1.

L4 ANSWER 38 OF 113 MEDLINE on STN

DUPLICATE 16

TI Onset of altered interventricular septal motion during cardiac surgery.
Assessment by continuous intraoperative transesophageal echocardiography.

L4 ANSWER 39 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

TI ECHOCARDIOGRAPHIC WALL MOTION ANALYSIS WITH 2-DIMENSIONAL FAST FOURIER
TRANSFORMATION.

L4 ANSWER 40 OF 113 MEDLINE on STN

DUPLICATE 17

TI Effects of morphine on gamma-aminobutyric acid turnover in the basal
ganglia. Possible correlation with its biphasic action on motility.

=> t ti l4 41-60

L4 ANSWER 41 OF 113 MEDLINE on STN

TI Echocardiographic documentation of the instant of cardiac rupture: a case
report.

L4 ANSWER 42 OF 113 MEDLINE on STN

DUPLICATE 18

TI Left ventricular **hyperkinesis** in acute myocardial infarction and
at control angiography after 1 month.

L4 ANSWER 43 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

TI CHILDHOOD DYSPHONIA.

L4 ANSWER 44 OF 113 MEDLINE on STN

TI [A rare cause of acute circulatory failure with pulmonary edema:
catecholergic cardiomyopathy of pheochromocytoma].
Une cause rare de defaillance circulatoire aigue avec oedeme pulmonaire:
la cardiomyopathie catecholergique du pheochromocytome.

L4 ANSWER 45 OF 113 MEDLINE on STN

DUPLICATE 19

TI Prolonged abnormalities of LV regional wall motion after normal
reperfusion in patients with preoperative cardiogenic shock.

L4 ANSWER 46 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS

RESERVED. on STN

TI What are the basal ganglia diseases?.

L4 ANSWER 47 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

TI PARKINSON'S SYNDROME PHENOMENA AFTER DELTA-SLEEP-INDUCING PEPTIDE
INJECTION INTO THE RAT'S SUBSTANTIA NIGRA.

L4 ANSWER 48 OF 113 MEDLINE on STN

DUPLICATE 20

TI Sequential noninvasive assessment of left ventricular size, regional wall
thickness and function during 3 hours of coronary artery occlusion and
reperfusion: differential effects of reflow in dogs with small vs large
areas at risk.

L4 ANSWER 49 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI THE FUNCTIONAL ANATOMY OF BASAL GANGLIA DISORDERS.

L4 ANSWER 50 OF 113 MEDLINE on STN DUPLICATE 21
TI Conditioning of morphine-induced locomotor activity and stereotyped behaviour in rats.

L4 ANSWER 51 OF 113 MEDLINE on STN DUPLICATE 22
TI [Myocardial asynergy of the left ventricle in patients with aortic stenosis].
Asinergii miokarda levogo zheludochka u bo'lnykh s stenozom us'tia aorty.

L4 ANSWER 52 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI DYSARTHRIAS OF MOVEMENT DISORDERS.

L4 ANSWER 53 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI DIAGNOSIS OF DYSKINESIAS OF THE GALLBLADDER BY ULTRASOUND TOMOGRAPHY BASIS OF AN EXTENDED INDICATION FOR CHOLECYSTECTOMY.

L4 ANSWER 54 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
DUPLICATE 23
TI DIAGNOSIS OF MOTOR DISORDERS OF THE DUODENUM.

L4 ANSWER 55 OF 113 MEDLINE on STN DUPLICATE 24
TI Effects of coronary artery surgery on left ventricular performance, segmental wall movement, and exertional ischaemia.

L4 ANSWER 56 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI PHYSICAL ACTIVITY REGIMEN IN PRESCHOOL CHILDREN AS A FUNCTION OF THE SEASON.

L4 ANSWER 57 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI AUGMENTATION OF CONTROL AND STABILIZATION IN NEURAL SYSTEMS.

L4 ANSWER 58 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI ECHOCARDIOGRAPHIC STUDY OF LEFT VENTRICULAR PERFORMANCE IN MITRAL VALVE PROLAPSE SYNDROME.

L4 ANSWER 59 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI QUANTITATIVE ANALYSIS OF REGIONAL WALL MOTION IN DOGS IN-SITU CHARACTERISTICS OF END-SYSTOLIC PRESSURE-LENGTH RELATION.

L4 ANSWER 60 OF 113 MEDLINE on STN DUPLICATE 25
TI Physiologic study of the terminal digestive tract in chronic painful constipation.

=> t ti 14 61-80

L4 ANSWER 61 OF 113 MEDLINE on STN
TI Mechanical and electrocardiographic sequence of coronary artery occlusion: an echocardiographic study during coronary angioplasty.

L4 ANSWER 62 OF 113 MEDLINE on STN DUPLICATE 26
TI Measurement of regional wall motion from biplane contrast ventriculograms: a comparison of the 30 degree right anterior oblique and 60 degree left anterior oblique projections in patients with acute myocardial infarction.

L4 ANSWER 63 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN
TI Fluvoxamine: A review of its pharmacodynamic and pharmacokinetic properties, and therapeutic efficacy in depressive illness.

L4 ANSWER 64 OF 113 MEDLINE on STN DUPLICATE 27
 TI [Mitral prolapse syndrome: clinical, electrocardiographic and
 angiocardiographic correlations. Study of 100 patients with healthy
 coronary vessels].
 Syndrome du prolapsus mitral: correlations clinique,
 electrocardiographique et angiographique. Etude de 100 patients a
 coronaires saines.

L4 ANSWER 65 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
 TI LEFT VENTRICULAR REGIONAL FUNCTION AND MYOCARDIAL METABOLIC CHANGES DURING
 ISCHEMIA.

L4 ANSWER 66 OF 113 MEDLINE on STN DUPLICATE 28
 TI Changes in left ventricular regional asynchrony after intracoronary
 thrombolysis in patients with impending myocardial infarction.

L4 ANSWER 67 OF 113 MEDLINE on STN DUPLICATE 29
 TI Asynchronous left ventricular wall motion early after coronary thrombosis.

L4 ANSWER 68 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
 RESERVED. on STN
 TI [Diagnosis of gallbladder dyskinesia by sonography during the stimulation
 of gallbladder].
 DIAGNOSTIK DER GALLENBLASENDYSKINESIEN MITTELS PHARMAKOSONOGRAPHIE.

L4 ANSWER 69 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
 RESERVED. on STN
 TI Three cases of lingual ballistic movement followed by rigid-dystonic state
 of the tongue.

L4 ANSWER 70 OF 113 MEDLINE on STN DUPLICATE 30
 TI Creatine kinase MB and M-mode echocardiographic changes in cardiac
 contusion.

L4 ANSWER 71 OF 113 MEDLINE on STN DUPLICATE 31
 TI Comparison of segmental and global ejection fraction in ischaemic heart
 disease.

L4 ANSWER 72 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
 DUPLICATE 32
 TI HYPERKINETIC CONTRACTION OF A NONISCHEMIC SEGMENT OF ISCHEMIC LEFT
 VENTRICLE IN ANESTHETIZED DOGS.

L4 ANSWER 73 OF 113 MEDLINE on STN DUPLICATE 33
 TI [Evaluation and identification of the extension of acute myocardial
 infarct and its complications by bidimensional echocardiography].
 Evaluacion e identificacion de la extension del infarto agudo al miocardio
 y sus complicaciones mediante la ecocardiografia bidimensional.

L4 ANSWER 74 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
 TI REST-EXERCISE CONTRAST VENTRICULOGRAPHIC WALL MOTION CHANGE POST
 THROMBOLYSIS.

L4 ANSWER 75 OF 113 MEDLINE on STN DUPLICATE 34
 TI Echocardiography in acute infectious myocarditis: relation to clinical and
 electrocardiographic findings.

L4 ANSWER 76 OF 113 MEDLINE on STN DUPLICATE 35
 TI [Effect of prolonged hypo- or **hyperkinesia** on the cAMP content
 in the liver, brain and muscles of rats].
 Vliianie na produlzhitelnata khipo- ili khiperkineziia vukhu sudurzhanieto
 na tsAMF v cherniia drob, mozuka i muskulite na plukhove.

L4 ANSWER 77 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN

TI ELECTRON MICROSCOPIC STUDY ON THE DIENCEPHALIC CHOROID PLEXUS IN RATS
AFTER PROLONGED **HYPOKINESIA** AND **HYPERKINESIA**.

L4 ANSWER 78 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI THE INFLUENCE OF THE MOTOR ACTIVITY REGIME ON THE ELABORATION CONSERVATION
AND REGENERATION OF MOTOR FOOD CONDITIONED REFLEXES IN RATS.

L4 ANSWER 79 OF 113 MEDLINE on STN DUPLICATE 36
TI Value of partial ejection fraction, volume increment, and regional wall
motion in identifying patients with clinically significant coronary artery
disease.

L4 ANSWER 80 OF 113 MEDLINE on STN DUPLICATE 37
TI Effect of interventions in salvaging left ventricular function in acute
myocardial infarction: a study of intracoronary streptokinase.

=> t ti l4 81-100

L4 ANSWER 81 OF 113 MEDLINE on STN DUPLICATE 38
TI Species differences in behavioural effects of rolipram and other adenosine
cyclic 3H, 5H-monophosphate phosphodiesterase inhibitors.

L4 ANSWER 82 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI How the initial vector reflects the left ventricular wall motion
abnormalities: Comparative study with vectorcardiographic and radionuclide
angiocardiographic findings.

L4 ANSWER 83 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Possible control of neuroleptic extrapyramidal symptomatology by lisuride.

L4 ANSWER 84 OF 113 MEDLINE on STN DUPLICATE 39
TI Evaluation of poor R wave progression by cross-sectional echocardiography
with wall motion index.

L4 ANSWER 85 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI NEURO CELLULAR POOL AND ADAPTATION IN THE INNERVATION SYSTEM OF ORGANS
DURING AGING AND IN HYPO KINESIA AND HYPER KINESIA.

L4 ANSWER 86 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI [Central motor and extrapyramidal side effects during therapy with
antidepressants].
ZENTRALMOTORISCHE UND EXTRAPYRAMIDALE NEBENWIRKUNGEN UNTER THERAPIE MIT
ANTIDEPRESSIVA.

L4 ANSWER 87 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI [Differentiation between diffuse myocardial disease and coronary heart
disease by cross sectional echocardiography].
DIFFERENTIALDIAGNOSE ZWISCHEN DIFFUSEN MYOKARDERKRANKUNGEN UND KORONARER
HERZKRANKHEIT MIT HILFE DER ZWEIDIMENSIONALEN ECHOKARDIOGRAPHIE.

L4 ANSWER 88 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI [The graphological pattern presented by the handwriting of cerebral
patients. Neurological graphology].
DAS SCHRIFTBILD BEI HIRNERKRANKUNGEN. NEUROLOGISCHE GRAPHOLOGIE.

L4 ANSWER 89 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI ASYNERGY OF LEFT VENTRICULAR CONTRACTIONS IN PATIENTS WITH ACQUIRED HEART
DEFECTS.

L4 ANSWER 90 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
DUPLICATE 40
TI NONINVASIVE EVALUATION OF LATE LEFT VENTRICULAR FUNCTION AFTER AORTIC
VALVE REPLACEMENT.

L4 ANSWER 91 OF 113 MEDLINE on STN
TI [Ultrastructural changes in the red bone marrow in hyper- and
hypokinesia].
Ultrastrukturni promeni v cherveniia kosten mozuk pri khiper- i
khipokineziia.

L4 ANSWER 92 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Minor physical anomalies in exceptional children.

L4 ANSWER 93 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Tardive dyskinesia.

L4 ANSWER 94 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Criteria and principles of operative tactics in children with
hydronephrosis.

L4 ANSWER 95 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
TI THE INTENSITY OF SYNTHESIS AND DECOMPOSITION OF TISSUE PROTEINS IN HYPO
KINESIA AND INCREASED MUSCULAR ACTIVITY.

L4 ANSWER 96 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
DUPLICATE 41
TI TOLERANCE TO HYPO KINESIA ELICITED BY DOPAMINE AGONISTS IN MICE HYPO
SENSITIZATION OF AUTO RECEPTORS.

L4 ANSWER 97 OF 113 MEDLINE on STN DUPLICATE 42
TI Ventricular buckling: a factor in the abnormal ventriculogram and peculiar
hemodynamics associated with mitral valve prolapse.

L4 ANSWER 98 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Ventricular buckling: a factor in the abnormal ventriculogram and peculiar
hemodynamics associated with mitral valve prolapse.

L4 ANSWER 99 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI [Brunkow's treatment technique from the neurosurgical point of view].
DIE BEHANDLUNGSTECHNIK BRUNKOW AUS DER SICHT DER NEUROCHIRURGIE.

L4 ANSWER 100 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Habitus of growing inbred animals under the conditions of hypo, normo and
hyperkinesia.

=> t ti l4 101-113

L4 ANSWER 101 OF 113 MEDLINE on STN DUPLICATE 43
TI [Primitive cardiomyopathies: dynamic geometry of left ventricular
contraction].
Miocardiopatie primitive: geometria dinamica della contrazione
ventricolare sinistra.

L4 ANSWER 102 OF 113 MEDLINE on STN DUPLICATE 44
TI Effect of CB 154 (2-bromo-alpha-ergocryptine) on paralysis agitans

compared with Madopar in a double-blind, cross-over trial.

- L4 ANSWER 103 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI [Treatment of Parkinson syndrome with levodopa plus carbidopa].
DIE BEHANDLUNG DES PARKINSONSCHEN SYNDROMS MIT NACOM.
- L4 ANSWER 104 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI [Drug induced extrapyramidal disturbances].
MEDIKAMENTOS BEDINGTE EXTRAPYRAMIDALE SYMPTOMATIK.
- L4 ANSWER 105 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Treatment of Parkinson's syndromes by a combination of L Dopa with
inhibitors of decarboxylase (Czech).
- L4 ANSWER 106 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Disruption of behavior in cats by chronic amphetamine intoxication.
- L4 ANSWER 107 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI The relation between the clinical effect of clozapine and the incidence of
side effects.
- L4 ANSWER 108 OF 113 MEDLINE on STN
TI [Changes in the large intestine after cholecystectomy].
Promeni v debeloto chervo sled kholetsistektomiia.
- L4 ANSWER 109 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI [Long term treatment of parkinsonism with levodopa plus a decarboxylase
inhibitor over a 3 year period].
LANGZEITTHERAPIE DES PARKINSONSYNDROMS MIT L DOPA UND EINEM
DECARBOXYLASEHEMMER UBER 3 JAHRE.
- L4 ANSWER 110 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Determination of the value of the Lombard test for the logopedic diagnosis
of phonatory disorders (Dutch).
- L4 ANSWER 111 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Movement disorders secondary to drugs.
- L4 ANSWER 112 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS
RESERVED. on STN
TI Survival time, Locomotor Activity (LA) and other parameters of starving
male and female mice.
- L4 ANSWER 113 OF 113 MEDLINE on STN
TI Morphologic correlates of azide-induced **hyperkinesis** and
hypokinesis.

=> d 14 bib,abs 22,93, 105, 107, 111

- L4 ANSWER 22 OF 113 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
AN 1994:41323 BIOSIS
DN PREV199497054323
TI Overview of basal ganglia function.
AU Delong, M. R.
CS Emory Univ. Sch. Med., Dep. Neurology, Suite 6000, 1639 Pierce Drive,

Atlanta, GA 30322, USA

SO Mano, N. [Editor]; Hamada, I. [Editor]; DeLong, M. R. [Editor]. Int. Congr. Ser. - Excerpta Med., (1993) pp. 65-70. International Congress Series; Role of the cerebellum and basal ganglia in voluntary movement. Publisher: Elsevier Science Publishers B.V., PO Box 211, Sara Burgerhartstraat 25, 1000 AE Amsterdam, Netherlands; Elsevier Science Publishing Co., Inc., P.O. Box 882, Madison Square Station, New York, New York 10159-2101, USA. Series: International Congress Series. Meeting Info.: 8th Tokyo Metropolitan Institute for Neuroscience (TMIN), International Symposium (20th Anniversary of TMIN). Tokyo, Japan. November 17-19, 1992. CODEN: EXMDA4. ISSN: 0531-5131. ISBN: 0-444-89813-1.

DT Book
Conference; (Meeting)
Book; (Book Chapter)

LA English

ED Entered STN: 3 Feb 1994
Last Updated on STN: 3 Feb 1994

L4 ANSWER 93 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN

AN 79243807 EMBASE

DN 1979243807

TI Tardive dyskinesia.

AU Gerlach J.

CS Dept. E., Sct. Hans Hosp., DK 4000, Denmark

SO Danish Medical Bulletin, (1979) 26/5 (209-245).
CODEN: DMBUAE

CY Denmark

DT Journal

FS 038 Adverse Reactions Titles
037 Drug Literature Index
032 Psychiatry
008 Neurology and Neurosurgery
019 Rehabilitation and Physical Medicine

LA English

AB Acute dystonia, parkinsonism and tardive dyskinesia (TD) are neurological side-effects in the traditional neuroleptic treatment of psychiatric patients. Persistent TD, in particular, is serious, since it indicates an irreversible neurotoxic effect. The actual research situation in the field of TD, particularly with regard to methodology and the actual results is reviewed. TD-like hyperkinetic movements are occasionally observed relatively early in neuroleptic treatment, possibly when the treatment is accentuated or on supplementary cholinergic treatment. This initial **hyperkinesia** is often, but not always, accompanied by dystonia, parkinsonism and/or akathisia, and decreases when the neuroleptic treatment is reduced and on anticholinergic treatment. Dystonia, parkinsonism and **hyperkinesia** may occur at any stage in the neuroleptic treatment, but dystonia, hypertonia and **hypokinesia** are particularly observed initially, while **hyperkinesia** dominates in the late treatment stage. The three syndromes thus overlap clinically, and have partly coincident pathogenetic mechanisms.

L4 ANSWER 105 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN

AN 77000717 EMBASE

DN 1977000717

TI Treatment of Parkinson's syndromes by a combination of L Dopa with inhibitors of decarboxylase (Czech).

AU Hanzal F.

CS Kat. Fak. Ped., KU, Praha, Czechoslovakia

SO Ceska a Slovenska Neurologie a Neurochirurgie, (1975) 38/6 (327-332).
CODEN: CKNNAS

DT Journal

FS 037 Drug Literature Index
 008 Neurology and Neurosurgery
 020 Gerontology and Geriatrics
 030 Pharmacology
 LA Czech
 AB Clinical research on the effect of a combination of L dopa with a decarboxylase inhibitor, carbidopa in 20 cases of parkinsonism between age 44 and 76 showed an average improvement of extrapyramidal signs of 53.3%. **Hypokinesia** improved on the average by 58%, rigidity by 56% and tremor by 46%. The daily maintenance dose is relatively low (2 to 3 tablets of L dopa at 250 mg and 25 mg carbidopa), producing no gastrointestinal symptoms in a specially tested group even after one year. Some unfavourable side effects at the beginning of treatment (oral automatism, **hyperkinesia**, paresthesia or extrasystoles) occurred in 3/4 of the cases, but cleared up in the course of 2-8 weeks with the exception of oral automatism in 2 patients. Combination of L dopa with decarboxylase inhibitors is effective even in the 20% of patients with intolerance of L dopa alone.

L4 ANSWER 107 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN
 AN 77030920 EMBASE
 DN 1977030920
 TI The relation between the clinical effect of clozapine and the incidence of side effects.
 AU Marecek P.; Faltus F.; Dolezalova V.
 CS Psychiat. Clin., Praha, Czechoslovakia
 SO Activitas Nervosa Superior, (1975) 17/4 (221-222).
 CODEN: ACNSAX
 DT Journal
 FS 038 Adverse Reactions Titles
 037 Drug Literature Index
 032 Psychiatry
 LA English

L4 ANSWER 111 OF 113 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED. on STN
 AN 74104471 EMBASE
 DN 1974104471
 TI Movement disorders secondary to drugs.
 AU Paulson G.W.
 CS Dept. Med., Ohio State Univ. Coll. Med., Columbus, Ohio, United States
 SO Ohio State Medical Journal, (1973) 69/9 (685-691).
 CODEN: OSMJAT
 DT Journal
 FS 038 Adverse Reactions Titles
 037 Drug Literature Index
 030 Pharmacology
 032 Psychiatry
 008 Neurology and Neurosurgery
 LA English
 AB Numerous drugs can produce a movement disorder as an unwanted side effect. In addition to drugs that produce **hypokinesia** or **hyperkinesia**, abnormal movements can result from toxins such as the heavy metals. The phenothiazines demonstrate the variety of effects since this group of drugs can produce, not only the well known 'pseudoparkinsonism,' but in addition, an acute dystonic reaction. A third effect is the 'tardive' or chronic dyskinesia seen after prolonged ingestion of phenothiazines. Tardive dyskinesia involves primarily hands and mouth, and may result for the first time when the medication is reduced in dosage. It was suggested that this movement disorder results from distortion in the receptor sites of the central nervous system. As uncertain as the explanation for this movement disorder, the explanation for the effects of drug withdrawal are even less well known. Whatever the

explanation for the tremulousness seen with withdrawal, development of a rational therapy for the phenomenon will represent a major therapeutic advance.

=> FIL STNGUIDE
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
31.42	31.63

FULL ESTIMATED COST

FILE 'STNGUIDE' ENTERED AT 14:28:43 ON 04 JUN 2004
USE IS SUBJECT TO THE TERMS OF YOUR CUSTOMER AGREEMENT.
COPYRIGHT (C) 2004 AMERICAN CHEMICAL SOCIETY, JAPAN SCIENCE
AND TECHNOLOGY CORPORATION, AND FACHINFORMATIONSZENTRUM KARLSRUHE

FILE CONTAINS CURRENT INFORMATION.
LAST RELOADED: May 28, 2004 (20040528/UP).

=> s dykinesa and dopamine and agonist and AMPA and antagonist

0 DYKINESA
0 DOPAMINE
0 AGONIST
0 AMPA
0 ANTAGONIST

L5 0 DYKINESA AND DOPAMINE AND AGONIST AND AMPA AND ANTAGONIST

=>

=>

=>

=> file medline biosis medline embase wpids
COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
0.48	32.11

FULL ESTIMATED COST

FILE 'MEDLINE' ENTERED AT 14:33:43 ON 04 JUN 2004

FILE 'BIOSIS' ENTERED AT 14:33:43 ON 04 JUN 2004
COPYRIGHT (C) 2004 BIOLOGICAL ABSTRACTS INC.(R)

FILE 'EMBASE' ENTERED AT 14:33:43 ON 04 JUN 2004
COPYRIGHT (C) 2004 Elsevier Inc. All rights reserved.

FILE 'WPIDS' ENTERED AT 14:33:43 ON 04 JUN 2004
COPYRIGHT (C) 2004 THOMSON DERWENT

=> s dykinesa and dopamine and agonist and AMPA and antagonist

L6 0 DYKINESA AND DOPAMINE AND AGONIST AND AMPA AND ANTAGONIST

=> s dykines? and dopamine and agonist and AMPA and antagonist

L7 0 DYKINES? AND DOPAMINE AND AGONIST AND AMPA AND ANTAGONIST

=> s dyskines? and dopamine and agonist and AMPA and antagonist

L8 10 DYSKINES? AND DOPAMINE AND AGONIST AND AMPA AND ANTAGONIST

=> dup rem

ENTER L# LIST OR (END):l8

PROCESSING COMPLETED FOR L8

L9 6 DUP REM L8 (4 DUPLICATES REMOVED)

=> d ti l9 1-6

L9 ANSWER 1 OF 6 MEDLINE on STN DUPLICATE 1
 TI A2A **antagonist** prevents **dopamine agonist**
 -induced motor complications in animal models of Parkinson's disease.

L9 ANSWER 2 OF 6 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.
 on STN
 TI Targeting striatal cholinergic interneurons in Parkinson's disease: Focus
 on metabotropic glutamate receptors.

L9 ANSWER 3 OF 6 MEDLINE on STN DUPLICATE 2
 TI Alteration of glutamate receptors in the striatum of dyskinetic
 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-treated monkeys following
dopamine agonist treatment.

L9 ANSWER 4 OF 6 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
 TI Methods of administering **AMPA** receptor antagonists to treat
dyskinesias associated with **dopamine agonist**
 therapy.

L9 ANSWER 5 OF 6 WPIDS COPYRIGHT 2004 THOMSON DERWENT on STN
 TI Use of **AMPA antagonist** for treatment of
dyskinesia associated with **dopamine agonist**
 therapy.

L9 ANSWER 6 OF 6 WPIDS COPYRIGHT 2004 THOMSON DERWENT on STN
 TI Treating **dyskinesia** associated with **dopamine**
agonist therapy in mammal - comprises administering **AMPA**
 receptor **antagonist**.

=> d bib abs 1-6

L9 ANSWER 1 OF 6 MEDLINE on STN DUPLICATE 1
 AN 2003557298 MEDLINE
 DN PubMed ID: 14637099
 TI A2A **antagonist** prevents **dopamine agonist**
 -induced motor complications in animal models of Parkinson's disease.
 CM Comment in: Exp Neurol. 2003 Nov;184(1):20-3. PubMed ID: 14637073
 AU Bibbiani F; Oh J D; Petzer J P; Castagnoli N Jr; Chen J-F; Schwarzschild M
 A; Chase T N
 CS ETB, NINDS, National Institutes of Health, Bethesda, MD 20892, USA.
 SO Experimental neurology, (2003 Nov) 184 (1) 285-94.
 Journal code: 0370712. ISSN: 0014-4886.
 CY United States
 DT Journal; Article; (JOURNAL ARTICLE)
 LA English
 FS Priority Journals
 EM 200401
 ED Entered STN: 20031126
 Last Updated on STN: 20040106
 Entered Medline: 20040105
 AB Adenosine A(2A) receptors, abundantly expressed on striatal medium spiny
 neurons, appear to activate signaling cascades implicated in the
 regulation of coexpressed ionotropic glutamatergic receptors. To evaluate
 the contribution of adenosinergic mechanisms to the pathogenesis of the
 response alterations induced by dopaminergic treatment, we studied the
 ability of the selective adenosine A(2A) receptor **antagonist**
 KW-6002 to prevent as well as palliate these syndromes in rodent and
 primate models of Parkinson's disease. In rats, KW-6002 reversed the
 shortened motor response produced by chronic levodopa treatment while
 reducing levodopa-induced hyperphosphorylation at S845 residues on
AMPA receptor GluR1 subunits. In primates, KW-6002 evidenced
 modest antiparkinsonian activity when given alone. Once-daily
 coadministration of KW-6002 with apomorphine prevented the development of

dyskinesias, which appeared in control animals 7-10 days after initiating apomorphine treatment. Animals initially given apomorphine plus KW-6002 for 3 weeks did not begin to manifest apomorphine-induced **dyskinesias** until 10-12 days after discontinuing the A(2A) **antagonist**. These results suggest that KW-6002 can attenuate the induction as well as the expression of motor response alterations to chronic dopaminergic stimulation in parkinsonian animals, possibly by blocking A(2A) receptor-stimulated signaling pathways. Our findings strengthen the rationale for developing A(2A) antagonists as an early treatment strategy for Parkinson's disease.

L9 ANSWER 2 OF 6 EMBASE COPYRIGHT 2004 ELSEVIER INC. ALL RIGHTS RESERVED.
on STN
AN 2003234243 EMBASE
TI Targeting striatal cholinergic interneurons in Parkinson's disease: Focus
on metabotropic glutamate receptors.
AU Pisani A.; Bonsi P.; Centonze D.; Gubellini P.; Bernardi G.; Calabresi P.
CS A. Pisani, Clinica Neurologica, Dipartimento di Neuroscienze, Univ. di
Roma Tor Vergata, Rome, Italy. pisani@uniroma2.it
SO Neuropharmacology, (2003) 45/1 (45-56).
Refs: 95
ISSN: 0028-3908 CODEN: NEPHBW
CY United Kingdom
DT Journal; (Short Survey)
FS 008 Neurology and Neurosurgery
030 Pharmacology
037 Drug Literature Index
038 Adverse Reactions Titles
LA English
SL English
AB In the early sixties, anticholinergic drugs were introduced in the
pharmacological treatment of Parkinson's disease (PD). The rationale
behind their utilisation in the treatment of the disease was based on the
evidence of an imbalance between the dopaminergic inputs and the intrinsic
cholinergic innervation within the striatum. Metabotropic glutamate (mGlu)
receptors have been shown to play a key role in striatal function both in
physiological conditions and in experimental models of diseases affecting
this brain area. Indeed, compelling electrophysiological and morphological
evidence shows that mGlu receptors are highly expressed at cellular level
and exert a profound modulatory role on cholinergic interneurons
excitability. This review will provide a brief survey of studies on the
localization and function of mGlu receptors in cholinergic interneurons.
The potential relevance of these findings in the control of motor function
and in the treatment of PD will be discussed. .COPYRGT. 2003 Elsevier
Science Ltd. All rights reserved.

L9 ANSWER 3 OF 6 MEDLINE on STN DUPLICATE 2
AN 2002119267 MEDLINE
DN PubMed ID: 11853103
TI Alteration of glutamate receptors in the striatum of dyskinetic
1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-treated monkeys following
dopamine agonist treatment.
AU Calon Frederic; Morissette Marc; Ghribi Othman; Goulet Martin; Grondin
Richard; Blanchet Pierre J; Bedard Paul J; Di Paolo Therese
CS Oncology and Molecular Endocrinology Research Center Laval University
Medical Center (CHUL), Quebec, Canada.
SO Progress in neuro-psychopharmacology & biological psychiatry, (2002 Jan)
26 (1) 127-38.
Journal code: 8211617. ISSN: 0278-5846.
CY England: United Kingdom
DT Journal; Article; (JOURNAL ARTICLE)
LA English
FS Priority Journals
EM 200208

ED Entered STN: 20020221
 Last Updated on STN: 20020821
 Entered Medline: 20020820

AB The effects of 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP)-induced nigrostriatal lesion and dopaminomimetic treatment on parameters of glutamatergic activity within the basal ganglia of monkeys were studied in relation with the development of **dyskinesias**. Drug-naive controls, saline-treated MPTP monkeys, as well as MPTP monkeys treated with either a long-acting D2 **agonist** (cabergoline) or a D1 **agonist** (SKF-82958) given by intermittent injections or continuous infusion, were included in this study. 3H-L-glutamate, 3H-alpha-amino-3-hydroxy-5-methylisoxasole-4-propionate (**AMPA**), 3H-glycine, 3H-CGP39653 (an N-methyl-D-aspartate, NMDA, **antagonist** selective for NR1/NR2A assembly) and 3H-Ro 25-6981 (an NMDA **antagonist** selective for NR1/NR2B assembly), specific binding to glutamate receptors, the expression of the NR1 subunit of NMDA receptors and glutamate, glutamine and glycine concentrations were studied by autoradiography, in situ hybridization and high-performance liquid chromatography (HPLC), respectively. Pulsatile SKF-82958 and cabergoline treatment relieved parkinsonian symptoms, whereas animals continuously treated with SKF-82958 remained akinetic. Pulsatile SKF-82958 induced **dyskinesias** in two of the three animals tested, whereas cabergoline did not. MPTP induced no significant changes of striatal specific binding of the radioligands used, NR1 mRNA expression and amino acid concentrations. In the putamen, pulsatile SKF-82958 treatment was associated with decreased content of glycine and glutamate, whereas only glycine was decreased in cabergoline-treated monkeys. Cabergoline and continuous administration of SKF-82958 led to lower levels of NR1 mRNA in the caudate in comparison to pulsatile SKF-82958 administration. The development of **dyskinesias** following a D1 **agonist** treatment was associated with an upregulation of 3H-glutamate [+49%], 3H-**AMPA** [+38%], 3H-CGP39653 [+ 111%], 3H-glycine [+ 26%, nonsignificant] and 3H-Ro 25-6981 [+ 33%] specific binding in the striatum in comparison to nondyskinetic MPTP monkeys. Our data suggest that supersensitivity to glutamatergic input in the striatum might play a role in the pathogenesis of dopaminomimetic-induced **dyskinesias** and further support the therapeutic potential of glutamate antagonists in Parkinson's disease.

L9 ANSWER 4 OF 6 BIOSIS COPYRIGHT 2004 BIOLOGICAL ABSTRACTS INC. on STN
 AN 2001:253193 BIOSIS
 DN PREV200100253193
 TI Methods of administering **AMPA** receptor antagonists to treat **dyskinesias** associated with **dopamine agonist** therapy.
 AU Chenard, Bertrand L. [Inventor]; Welch, Willard M. [Inventor]; Menniti, Frank S. [Inventor, Reprint author]
 CS Mystic, CT, USA
 ASSIGNEE: Pfizer Inc
 PI US 6136812 October 24, 2000
 SO Official Gazette of the United States Patent and Trademark Office Patents, (Oct. 24, 2000) Vol. 1239, No. 4. e-file.
 CODEN: OGUPE7. ISSN: 0098-1133.
 DT Patent
 LA English
 ED Entered STN: 23 May 2001
 Last Updated on STN: 19 Feb 2002

AB The invention relates to a method of treating **dyskinesias** associated with **dopamine agonist** therapy in a mammal which comprises administering to said mammal a compound, as defined herein, which is an **antagonist** of the **AMPA** receptor. **Dopamine agonist** therapy, as referred to in the present invention, is generally used in the treatment of a central nervous system disorder such as Parkinson's disease.

L9 ANSWER 5 OF 6 WPIDS COPYRIGHT 2004 THOMSON DERWENT on STN
 AN 1999-245888 [21] WPIDS
 CR 1999-169071 [15]
 DNC C1999-071960
 TI Use of **AMPA antagonist** for treatment of
dyskinesia associated with **dopamine agonist**
 therapy.
 DC B02 B05
 IN CHENARD, B L; MENNITI, F S; WELCH, W M; WELCH, W K; MCKOWAN, W W
 PA (PFIZ) PFIZER PROD INC; (PFIZ) PFIZER INC
 CYC 33
 PI EP 900568 A2 19990310 (199921)* EN 22
 R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT
 RO SE SI
 AU 9883120 A 19990318 (199923)
 HU 9802021 A2 19990528 (199930)
 CA 2246839 A 19990305 (199933)
 JP 11158072 A 19990615 (199934) 43
 KR 99029570 A 19990426 (200028)
 ZA 9808139 A 20000531 (200032) 440
 NZ 331741 A 20000825 (200049)
 US 6136812 A 20001024 (200055)
 AU 736254 B 20010726 (200149)
 JP 2001316267 A 20011113 (200207) 36
 CA 2246839 C 20021112 (200302) EN
 ADT EP 900568 A2 EP 1998-307181 19980904; AU 9883120 A AU 1998-83120 19980904;
 HU 9802021 A2 HU 1998-2021 19980904; CA 2246839 A CA 1998-2246839
 19980908; JP 11158072 A JP 1998-245269 19980831; KR 99029570 A KR
 1998-36653 19980905; ZA 9808139 A ZA 1998-8139 19980907; NZ 331741 A NZ
 1998-331741 19980904; US 6136812 A Provisional US 1997-58098P 19970905, US
 1998-148974 19980904; AU 736254 B AU 1998-83120 19980904; JP 2001316267 A
 Div ex JP 1998-245269 19980831, JP 2001-134816 19980831; CA 2246839 C CA
 1998-2246839 19980908
 FDT AU 736254 B Previous Publ. AU 9883120
 PRAI US 1997-58098P 19970905; US 1997-57987P 19970905;
 US 1998-148974 19980904
 AN 1999-245888 [21] WPIDS
 CR 1999-169071 [15]
 AB EP 900568 A UPAB: 20030111
 NOVELTY - Use of an alpha -amino-3-hydroxy-5-methyl-4-isoxazolepropionic
 acid (**AMPA**) **antagonist** for treatment of
dyskinesia associated with **dopamine agonist**
 therapy is new
 DETAILED DESCRIPTION - Use of an alpha -amino-3-hydroxy-5-methyl-4-
 isoxazolepropionic acid (**AMPA**) **antagonist** known from
 WO9743276, PCT/IB9800150, EP98304319.0, EP98304522.0, PCT/IB98/00151 or US
 provisional number 60/057990 for treatment of **dyskinesia**
 associated with **dopamine agonist** therapy is new. The
 compound is e.g. selected from a list of over 200 specific compounds e.g.:
 (1) (S)-3-(2-chlorophenyl)-2-(2-(6-diethylaminomethyl-pyridin-2-yl)-
 vinyl)-6-fluoro-3H-quinazolin-4-one;
 (2) (S)-2-(2-(6-fluoro-3-(2-methyl-pyridin-3-yl)-4-oxo-3,4-dihydro-
 quinazolin-2-yl)-vinyl)-benzonitrile;
 (3) 3-(2-chlorophenyl)-2-(2-pyridin-2-yl-vinyl)-3H-quinazolin-4-one;
 (4) 6-chloro-3-(2-chlorophenyl)-2-(2-hydroxy-2-(6-methylpyridin-2-
 yl)vinyl)-3H-quinazolin-4-one; or
 (5) 3-(2-chlorophenyl)-6-fluoro-2-((6-methyl-pyridin-2-
 ylaminomethyl)-3H-quinazolin-4-one.
 ACTIVITY - None given.
 MECHANISM OF ACTION - **AMPA antagonist**. The
 compounds are stated to inhibit **AMPA** receptor activation induced
 45Ca2+ uptake by 50 % or more at 0.5 micro M (no specific data given).
 USE - The method is used to treat **dyskinesias** caused by

administration of L-dopa optionally in combination with a peripheral dopadecarboxylase inhibitor (especially carbidopa or benserazide) for the treatment of Parkinson's disease (claimed). **Dyskinesias** treated include chorea, tremor, ballism, dystonia, athetosis, myoclonus or tic.

A test to assess the efficacy of the compounds against **dyskinesias** is described, but no results are given.
Dwg.0/0

L9 ANSWER 6 OF 6 WPIDS COPYRIGHT 2004 THOMSON DERWENT on STN

AN 1999-155778 [14] WPIDS

DNC C1999-046029

TI Treating **dyskinesia** associated with **dopamine** **agonist** therapy in mammal - comprises administering **AMPA** receptor **antagonist**.

DC B02 B05

IN CHENARD, B L; GREENAMYRE, J T; MENNITI, F S; WELCH, W M; MCKOWAN, W W
PA (GREE-I) GREENAMYRE J; (PFIZ) PFIZER PROD INC; (GREE-I) GREENAMYRE J T;
(GREE-I) GREENAMAYER J T; (CHEN-I) CHENARD B L; (MENN-I) MENNITI F S;
(WELC-I) WELCH W M

CYC 34

PI EP 900567 A2 19990310 (199914)* EN 6
R: AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL PT
RO SE SI

AU 9883193 A 19990318 (199923)
HU 9802022 A2 19990528 (199930)
JP 11139991 A 19990525 (199931) 7
CA 2246560 A 19990305 (199933)
KR 99029528 A 19990426 (200028)
ZA 9808009 A 20000531 (200032) 13
NZ 331636 A 20000825 (200049)
US 2001034345 A1 20011025 (200170)
CA 2246560 C 20021217 (200309) EN
TW 490304 A 20020611 (200321)
AU 2002300534 A1 20030213 (200427)#

ADT EP 900567 A2 EP 1998-306661 19980820; AU 9883193 A AU 1998-83193 19980907;
HU 9802022 A2 HU 1998-2022 19980904; JP 11139991 A JP 1998-249644
19980903; CA 2246560 A CA 1998-2246560 19980903; KR 99029528 A KR
1998-36442 19980904; ZA 9808009 A ZA 1998-8009 19980902; NZ 331636 A NZ
1998-331636 19980831; US 2001034345 A1 Provisional US 1997-57965P
19970905, US 1998-148973 19980904; CA 2246560 C CA 1998-2246560 19980903;
TW 490304 A TW 1998-114576 19980902; AU 2002300534 A1 Div ex AU 1998-83193
19980907, AU 2002-300534 20020814

PRAI US 1997-57965P 19970905; US 1998-148973 19980904;
AU 2002-300534 20020814

AN 1999-155778 [14] WPIDS

AB EP 900567 A UPAB: 19990412
NOVELTY - The method comprises administering to the mammal an **AMPA** receptor **antagonist**.

USE - The method is useful for treating **dyskinesias** in mammals.

ADVANTAGE - The method effectively treats **dyskinesia**.

Dwg.0/0

=> logoff hold

COST IN U.S. DOLLARS

SINCE FILE	TOTAL
ENTRY	SESSION
18.53	50.64

FULL ESTIMATED COST

SESSION WILL BE HELD FOR 60 MINUTES

STN INTERNATIONAL SESSION SUSPENDED AT 14:36:45 ON 04 JUN 2004

Connecting via Winsock to STN

Welcome to STN International! Enter x:x

LOGINID:SSSPTA1639MLS

PASSWORD:

TERMINAL (ENTER 1, 2, 3, OR ?):2

* * * * * Welcome to STN International * * * * *

NEWS 1		Web Page URLs for STN Seminar Schedule - N. America
NEWS 2		"Ask CAS" for self-help around the clock
NEWS 3	JAN 27	Source of Registration (SR) information in REGISTRY updated and searchable
NEWS 4	JAN 27	A new search aid, the Company Name Thesaurus, available in CA/CAPLUS
NEWS 5	FEB 05	German (DE) application and patent publication number format changes
NEWS 6	MAR 03	MEDLINE and LMEADLINE reloaded
NEWS 7	MAR 03	MEDLINE file segment of TOXCENTER reloaded
NEWS 8	MAR 03	FRANCEPAT now available on STN
NEWS 9	MAR 29	Pharmaceutical Substances (PS) now available on STN
NEWS 10	MAR 29	WPIFV now available on STN
NEWS 11	MAR 29	New monthly current-awareness alert (SDI) frequency in RAPRA
NEWS 12	APR 26	PROMT: New display field available
NEWS 13	APR 26	IFIPAT/IFIUDB/IFICDB: New super search and display field available
NEWS 14	APR 26	LITALERT now available on STN
NEWS 15	APR 27	NLDB: New search and display fields available
NEWS 16	May 10	PROUSDDR now available on STN
NEWS 17	May 19	PROUSDDR: One FREE connect hour, per account, in both May and June 2004
NEWS 18	May 12	EXTEND option available in structure searching
NEWS 19	May 12	Polymer links for the POLYLINK command completed in REGISTRY
NEWS 20	May 17	FRFULL now available on STN
NEWS 21	May 27	STN User Update to be held June 7 and June 8 at the SLA 2004 Conference
NEWS 22	May 27	New UPM (Update Code Maximum) field for more efficient patent SDIs in CAPLUS
NEWS 23	May 27	CAPLUS super roles and document types searchable in REGISTRY
NEWS 24	May 27	Explore APOLLIT with free connect time in June 2004
NEWS EXPRESS		MARCH 31 CURRENT WINDOWS VERSION IS V7.00A, CURRENT MACINTOSH VERSION IS V6.0c(ENG) AND V6.0Jc(JP), AND CURRENT DISCOVER FILE IS DATED 26 APRIL 2004
NEWS HOURS		STN Operating Hours Plus Help Desk Availability
NEWS INTER		General Internet Information
NEWS LOGIN		Welcome Banner and News Items
NEWS PHONE		Direct Dial and Telecommunication Network Access to STN
NEWS WWW		CAS World Wide Web Site (general information)

Enter NEWS followed by the item number or name to see news on that specific topic.

All use of STN is subject to the provisions of the STN Customer agreement. Please note that this agreement limits use to scientific research. Use for software development or design or implementation of commercial gateways or other similar uses is prohibited and may